

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

STAMICIS 1 mg kit for radiopharmaceutical preparation

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 1 mg [Tetrakis (2-methoxyisobutyl isonitrile) copper (I)] tetrafluoroborate.

The radionuclide is not part of the kit.

Excipients with known effect:

One ml of solution contains 4.5 mg of sodium

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Kit for radiopharmaceutical preparation.

White powder.

### 4. CLINICAL PARTICULARS

#### 4.1. Therapeutic indications

This medicinal product is for diagnostic use only.

This is indicated for adults. For paediatric population see section 4.2.

After radiolabelling with sodium pertechnetate ( $^{99m}\text{Tc}$ ) solution, the solution of technetium ( $^{99m}\text{Tc}$ ) sestamibi obtained is indicated for:

- **Myocardial perfusion scintigraphy** for the detection and localisation of coronary artery disease (angina pectoris and myocardial infarction).
- **Assessment of global ventricular function.** First-pass technique for determination of ejection fraction and/or ECG-triggered, gated SPECT for evaluation of left ventricular ejection fraction, volumes and regional wall motion.
- **Scintimammography for the detection of suspected breast cancer** when mammography is equivocal, inadequate or indeterminate.
- **Localisation of hyperfunctioning parathyroid tissue** in patients with recurrent or persistent disease in both primary and secondary hyperparathyroidism, and in patients with primary hyperparathyroidism scheduled to undergo initial surgery of the parathyroid glands.

## 4.2 Posology and method of administration

### Posology

#### Adults and elderly population

Posology may vary depending on gamma camera characteristics and reconstruction modalities. The injection of activities greater than local DRLs (Diagnostic Reference Levels) should be justified.

The recommended activity range for intravenous administration to an adult patient of average weight (70 kg) is for:

#### *Diagnosis of reduced coronary perfusion and myocardial infarction*

The recommended activity range for diagnosis of ischaemic heart disease according to the European Nuclear Medicine Guide of the EANM 2019 (European Association of Nuclear Medicine) is:

- Two-day protocol: 300-600 MBq/study if a conventional Anger gamma camera is used, or 180-500 MBq if a cardiac-centered gamma camera is used for acquisition
- One-day protocol: 250-400 MBq for the first injection, three times more for the second injection if a conventional Anger gamma camera is used, or 150-300 MBq if a cardiac-centered gamma camera is used for acquisition.

Not more than a total of 1,600 MBq should be administered for a one-day protocol and 1,200 MBq for a two-day-protocol. For a one-day protocol, the two injections (stress and rest) should be done at least two hours apart but may be performed in either order. After the stress injection, exercise should be encouraged for an additional one minute (if possible).

For diagnosis of myocardial infarction one injection at rest is usually sufficient.

For diagnosis of ischaemic heart disease two injections (stress and rest) are required in order to differentiate transiently from persistently reduced myocardial uptake.

*Assessment of global ventricular function:* 600-800 MBq injected as a bolus.

*Scintimammography:* 700-1000 MBq injected as a bolus usually in the arm opposite to the lesion.

#### *Localisation of hyperfunctioning parathyroid tissue:*

For the dual phase technique 400-900 MBq injected as a bolus. The typical activity is between 500-700 MBq.

For the subtraction technique with sodium pertechnetate ( $^{99m}\text{Tc}$ ): 75-110 MBq sodium pertechnetate ( $^{99m}\text{Tc}$ ) by intravenous administration followed by 400-900 MBq ( $^{99m}\text{Tc}$ ) sestamibi by intravenous administration, or 400-900 MBq ( $^{99m}\text{Tc}$ ) sestamibi by intravenous administration followed by 150 MBq sodium pertechnetate ( $^{99m}\text{Tc}$ ) by intravenous administration.

For the subtraction technique with sodium iodide ( $^{123}\text{I}$ ): 7.5-15 MBq sodium iodide ( $^{123}\text{I}$ ) administered orally or intravenously followed 2 hours later by 400-900 MBq ( $^{99m}\text{Tc}$ ) sestamibi by intravenous administration.

#### Renal impairment

Careful consideration of the activity to be administered is required since an increased radiation exposure is possible in these patients.

#### Hepatic impairment

In general, activity selection for patients with a decreased hepatic function should be cautious, usually starting at the low end of the dosing range.

### Paediatric population

The use in children and adolescents has to be considered carefully, based upon clinical needs and assessing the risk/benefit ratio in this patient group. The activities to be administered to children and adolescents may be calculated according to the recommendations of the European Association of Nuclear Medicine (EANM) paediatric dosage card; the activity administered to children and to adolescents may be calculated by multiplying a baseline activity (for calculation purposes) by the weight-dependent multiples given in the table below.

$A[\text{MBq}]_{\text{Administered}} = \text{Baseline Activity} \times \text{Multiple}$ :

The baseline activity is 63 MBq as a cancer seeking agent. For cardiac imaging, the minimum and maximum baseline activities are 42 and 63 MBq, respectively, for the two-day protocol cardiac scan both at rest and stress. For the one-day cardiac imaging protocol, the baseline activity is 28 MBq at rest and 84 MBq at stress. The minimum activity for any imaging study is 80 MBq.

<b>Weight [kg]</b>	<b>Multiple</b>	<b>Weight [kg]</b>	<b>Multiple</b>	<b>Weight [kg]</b>	<b>Multiple</b>
<b>3</b>	1	<b>22</b>	5.29	<b>42</b>	9.14
<b>4</b>	1.14	<b>24</b>	5.71	<b>44</b>	9.57
<b>6</b>	1.71	<b>26</b>	6.14	<b>46</b>	10.00
<b>8</b>	2.14	<b>28</b>	6.43	<b>48</b>	10.29
<b>10</b>	2.71	<b>30</b>	6.86	<b>50</b>	10.71
<b>12</b>	3.14	<b>32</b>	7.29	<b>52-54</b>	11.29
<b>14</b>	3.57	<b>34</b>	7.72	<b>56-58</b>	12.00
<b>16</b>	4.00	<b>36</b>	8.00	<b>60-62</b>	12.71
<b>18</b>	4.43	<b>38</b>	8.43	<b>64-66</b>	13.43
<b>20</b>	4.86	<b>40</b>	8.86	<b>68</b>	14.00

### Method of administration

For intravenous use.

Because of potential tissue damage, extravasal injection of this radioactive product has to be strictly avoided.

For multidose use.

### *Precautions to be taken before handling or administration of the medicinal product*

This medicinal product should be reconstituted before administration to the patient. For instructions on reconstitution and control of the radiochemical purity of the medicinal product before administration, see section 12.

For patient preparation, see section 4.4.

### Image acquisition

#### *Cardiac imaging*

Imaging should begin approximately after 30-60 min after injection to allow for hepatobiliary clearance. Longer delay can be required for resting images and for stress with vasodilators alone because of the risk of higher subdiaphragmatic technetium ( $^{99\text{m}}\text{Tc}$ ) activity. There is no evidence for significant changes in myocardial tracer concentration or redistribution, therefore imaging for up to 6 hours post injection is possible. Test may be done in a one day or two days protocol.

Preferably tomographic imaging (SPECT) with or without ECG gating should be performed.

### *Scintimammography*

Breast imaging is optimally initiated 5 to 10 minutes post injection with the patient in the prone position with breast freely pendant.

The product is administered in an arm vein contralateral to the breast with the suspected abnormality. If the disease is bilateral, the injection is ideally administered in a dorsal vein of the foot.

#### Conventional gamma camera

The patient should then be repositioned so that the contralateral breast is pendant and a lateral image of it should be obtained. An anterior supine image may then be obtained with the patient's arms behind her head.

#### Camera dedicated to breast imaging

In case a camera dedicated to breast imaging is used, a relevant machine-specific protocol must be followed to obtain the best possible imaging performance.

### Parathyroid imaging

Parathyroid image acquisition depends on the protocol chosen. The most used studies are either the subtraction and/or the dual-phase techniques, which can be performed together.

For the subtraction technique either sodium iodide ( $^{123}\text{I}$ ) or sodium pertechnetate ( $^{99\text{m}}\text{Tc}$ ) can be used for imaging for the thyroid gland since these radiopharmaceuticals are trapped by functioning thyroid tissue. This image is subtracted from the technetium ( $^{99\text{m}}\text{Tc}$ ) sestamibi image, and pathological hyperfunctioning parathyroid tissue remains visible after subtraction.

When sodium iodide ( $^{123}\text{I}$ ) is used, images are acquired simultaneously, starting 5 min after injection of ( $^{99\text{m}}\text{Tc}$ )sestamibi. Images are inspected visually, normalized to thyroid counts, and sodium iodide ( $^{123}\text{I}$ ) images are subtracted from the ( $^{99\text{m}}\text{Tc}$ )sestamibi images.

When sodium pertechnetate ( $^{99\text{m}}\text{Tc}$ ) is used, image acquisition of sodium pertechnetate ( $^{99\text{m}}\text{Tc}$ ) is starting 20-30 minutes after injection. Image acquisition of ( $^{99\text{m}}\text{Tc}$ )sestamibi is starting 10-15 minutes after injection. Images of sodium pertechnetate ( $^{99\text{m}}\text{Tc}$ ) are either digitally or cognitively subtracted from the ( $^{99\text{m}}\text{Tc}$ ) sestamibi images.

When the dual phase technique is used, the first neck and mediastinum image is obtained 10 minutes later. After a wash-out period of 1 to 2 hours, neck and mediastinum imaging is again performed.

The planar images may be complemented by early and delayed SPECT or SPECT/CT.

### **4.3 Contraindications**

Hypersensitivity to the active substances to any of the excipients listed in section 6.1 or to any of the components of the labelled radiopharmaceutical.

In myocardial scintigraphy investigations under stress conditions, the general contraindications associated with the induction of ergometric or pharmacological stress should be considered.

### **4.4 Special warnings and precautions for use**

#### Potential for hypersensitivity or anaphylactic reactions

If hypersensitivity or anaphylactic reactions occur, the administration of the medicinal product must be discontinued immediately and intravenous treatment initiated, if necessary. To enable immediate action in emergencies, the necessary medicinal products and equipment such as endotracheal tube and ventilator must be immediately available.

#### Individual benefit/risk justification

For each patient, the radiation exposure must be justifiable by the likely benefit. The activity administered should in every case be as low as reasonably achievable to obtain the required diagnostic information.

#### Renal or hepatic impairment

Careful consideration of the benefit risk ratio in these patients is required since an increased radiation exposure is possible (see section 4.2).

#### Paediatric population

For information on the use in paediatric population, see section 4.2.

Careful consideration of the indication is required since the effective dose per MBq is higher than in adults (see section 11).

#### Patient preparation

The patient should be well hydrated before the start of the examination and urged to void as often as possible during the first hours after the examination in order to reduce radiation.

#### Cardiac imaging

If possible, patients should fast for at least four hours prior to the study. It is recommended that patients eat a light fatty meal or drink a glass or two of milk after each injection, prior to imaging. This will promote rapid hepatobiliary clearance of technetium ( $^{99m}\text{Tc}$ ) sestamibi resulting in less liver activity in the image.

#### Interpretation of technetium ( $^{99m}\text{Tc}$ ) sestamibi images

##### Interpretation of scintimammography

Breast lesions less than 1 cm in diameter may not all be detected with scintimammography as the sensitivity of technetium ( $^{99m}\text{Tc}$ ) sestamibi for the detection of these lesions is low. A negative examination does not exclude breast cancer especially in such a small lesion.

##### After the procedure

Close contact with infants and pregnant women should be restricted during the initial 24 hours following the injection.

##### Specific warnings

In myocardial scintigraphy investigations under stress conditions, the general contraindications and precautions associated with the induction of ergometric or pharmacological stress should be considered.

This medicinal product contains less than 1 mmol sodium (23 mg) per vial, i.e. essentially 'sodium-free'.

For precautions with respect to environmental hazard see section 6.6.

## **4.5 Interaction with other medicinal products and other forms of interaction**

#### Cardiac medications

Medicinal products which affect myocardial function and/or blood flow may cause false negative results in the diagnosis of coronary arterial disease. Particularly beta-blockers and calcium antagonists reduce oxygen consumption and thus also affect perfusion and beta-blockers inhibit the increase of heart frequency and blood pressure under stress. For this reason, concomitant medicinal product should be taken into consideration when interpreting the results of the scintigraphic examination. The recommendations of the applicable guidelines on ergometric or pharmacological stress tests should be followed.

### Proton pump inhibitors

The use of proton pump inhibitors has shown to be significantly associated with gastric wall uptake. Its proximity to the inferior myocardial wall may lead to either false-negative or false-positive findings, and therefore to an inaccurate diagnosis. A withdrawal period for at least 3 days is recommended.

### Iodine products

When the subtraction technique is used for imaging of hyperfunctioning parathyroid tissue, recent use of iodine containing radiologic contrast media, medicinal products used to treat hyper- or hypothyroidism or of several other medicinal products is likely to decrease the quality of thyroid imaging and even makes subtraction impossible. For a complete list of possibly interacting medicinal products refer to the SmPCs of sodium iodide (<sup>123</sup>I) or sodium pertechnetate (<sup>99m</sup>Tc).

### Paediatric population

Interaction studies have only been performed in adults.

## **4.6 Fertility, pregnancy and lactation**

### Women of childbearing potential

When an administration of radiopharmaceuticals to a woman of childbearing potential is intended, it is important to determine whether or not she is pregnant. Any woman who has missed a period should be assumed to be pregnant until proven otherwise. If in doubt about her potential pregnancy (if the woman has missed a period, if the period is very irregular, etc.), alternative techniques not using ionising radiation (if there are any) should be offered to the patient.

### Pregnancy

Radionuclide procedures carried out on pregnant women also involve radiation dose to the foetus. Only essential investigations should therefore be carried out during pregnancy, when the likely benefit far exceeds the risk incurred by the mother and foetus.

### Breastfeeding

Before administering radiopharmaceuticals to a mother who is breastfeeding consideration should be given to the possibility of delaying the administration of radionuclide until the mother has ceased breastfeeding, and to what is the most appropriate choice of radiopharmaceuticals, bearing in mind the secretion of activity in breast milk. If the administration is considered necessary, breastfeeding should be interrupted for 24 hours and the expressed feeds discarded.

Close contact with infants should be restricted during the initial 24 hours following injection.

### Fertility

No studies on fertility have been performed.

## **4.7 Effects on ability to drive and use machines**

STAMICIS has no or negligible influence on the ability to drive and use machines.

## **4.8 Undesirable effects**

The following table presents how the frequencies are reflected in this section:

Very common ( $\geq 1/10$ )
Common ( $\geq 1/100$ to $< 1/10$ )
Uncommon ( $\geq 1/1,000$ to $< 1/100$ )
Rare ( $\geq 1/10,000$ to $< 1/1,000$ )
Very rare ( $< 1/10,000$ )
Not known (cannot be estimated from the available data)

*Immune system disorders:*

Rare: Severe hypersensitivity reactions such as dyspnoea, hypotension, bradycardia, asthenia and vomiting (usually within two hours of administration), angioedema. Other hypersensitivity reactions (allergic skin and mucosa reactions with exanthema (pruritus, urticaria, oedema), vasodilatation).

Very rare: Other hypersensitivity reactions have been described in predisposed patients.

*Nervous system disorders:*

Uncommon: Headache.

Rare: Seizures (shortly after administration), syncope.

*Cardiac disorders*

Uncommon: Chest pain/angina pectoris, abnormal ECG.

Rare: Arrhythmia.

*Gastrointestinal disorders:*

Uncommon: Nausea.

Rare: Abdominal pain.

*Skin and subcutaneous tissue disorders:*

Rare: local reactions at the injection site, hypoaesthesia and paraesthesia, flushing.

Not known: Erythema multiforme.

*General disorders and administration site conditions:*

Common: Immediately after injection, a metallic or bitter taste, partly in combination with dry mouth and an alteration in the sense of smell may be observed.

Rare: Fever, fatigue, dizziness, transient arthritic-like pain.

*Other disorders*

Exposure to ionising radiation is linked with cancer induction and a potential for development of hereditary defects. As the effective dose is about 13.0 mSv when the maximal recommended activity of 1,600 MBq (400 at rest and 1,200 MBq at stress) for a 1-day-protocol is administered these adverse reactions are expected to occur with a low probability.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

## **4.9 Overdose**

In the event of administration of a radiation overdose with technetium (<sup>99m</sup>Tc) sestamibi the absorbed dose to the patient should be reduced where possible by increasing the elimination of the radionuclide from the body by frequent micturition and defaecation. It might be helpful to estimate the effective dose that was applied.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**


Pharmacotherapeutic group : diagnostic radiopharmaceuticals, Technetium (<sup>99m</sup>Tc) compounds, ATC code : V09GA01

### Pharmacodynamic effects

At the chemical concentrations used for diagnostic examinations, technetium ( $^{99m}\text{Tc}$ ) sestamibi solution does not appear to have any pharmacodynamic activity.

## **5.2 Pharmacokinetic properties**

After reconstitution with sodium pertechnetate ( $^{99m}\text{Tc}$ ), the following technetium ( $^{99m}\text{Tc}$ ) sestamibi complex is formed:



### Biodistribution

Technetium ( $^{99m}\text{Tc}$ ) sestamibi from the blood is rapidly distributed into the tissue: 5 minutes after injection only about 8% of the injected dose remains in the blood pool. In physiological distribution, evident concentration of technetium ( $^{99m}\text{Tc}$ ) sestamibi can be seen in vivo in several organs. In particular, normal tracer uptake is evident in the salivary glands, thyroid, myocardium, liver, gallbladder, small and large intestine, kidneys, bladder, choroid plexuses and skeletal muscles, occasionally in the nipples. Faint homogeneous uptake in the breast or axilla is normal.

### Myocardial perfusion scintigraphy

Technetium ( $^{99m}\text{Tc}$ ) sestamibi is a cationic complex which diffuses passively through the capillary and cell membrane. Within the cell it is localised in the mitochondria, where it is trapped, and retention is based on intact mitochondria, reflecting viable myocytes. After intravenous injection, it is distributed within the myocardium according to myocardial perfusion and viability. Myocardial uptake which is coronary flow dependent is 1.5% of the injected dose at stress and 1.2% of the injected dose at rest. Irreversibly damaged cells however do not take up technetium ( $^{99m}\text{Tc}$ ) sestamibi. The myocardial extraction level is reduced by hypoxia. It has very little redistribution and so separate injections are required for stress and resting studies.

### Scintimammography

The tissue uptake of technetium ( $^{99m}\text{Tc}$ ) sestamibi depends primarily on the vascularisation which is generally increased in tumor tissue. Technetium ( $^{99m}\text{Tc}$ ) sestamibi accumulates in various neoplasms and most markedly in mitochondria. Its uptake is related to increased energy-dependent metabolism and cell proliferation. Its cellular accumulation is reduced when multidrug resistance proteins are overexpressed.

### Parathyroid imaging of hyperfunctioning tissue

Technetium ( $^{99m}\text{Tc}$ ) sestamibi localises in both parathyroid tissue and functioning thyroid tissue but usually washes out of normal thyroid tissue more rapidly than out of abnormal parathyroid tissue.

### Elimination

Elimination of technetium ( $^{99m}\text{Tc}$ ) sestamibi occurs mostly through the kidneys and the hepatobiliary system. Activity of technetium ( $^{99m}\text{Tc}$ ) sestamibi from the gallbladder appears in the intestine within one hour of injection. About 27% of the injected dose is cleared through renal elimination after 24 hours and approximately 33% of the injected dose is cleared through the faeces in 48 hours. The pharmacokinetics in patients with renal or hepatic impairment has not been characterised.

### Half-Life

The biological myocardial half-life of technetium ( $^{99m}\text{Tc}$ ) sestamibi is approximately 7 hours at rest and stress. The effective half-life (which includes biological and physical half-lives) is approximately 3 hours for the heart and approximately 30 minutes for the liver.



### **5.3 Preclinical safety data**

In acute intravenous toxicity studies in mice, rats and dogs, the lowest dose of the reconstituted kit that resulted in any deaths was 7 mg/kg (expressed as Cu (MIBI)<sub>4</sub> BF<sub>4</sub> content) in female rats. This corresponds to 500 times the maximal human dose (MHD) of 0.014 mg/kg for adults (70 kg).

Neither rats nor dogs exhibited treatment related effects at reconstituted kit doses of 0.42 mg/kg (30 times MHD) and 0.07 mg/kg (5 times MHD) respectively for 28 days. At repeated dose administration, the first toxicity symptoms appeared during the administration of 150 times the daily dose during 28 days.

Extravasation administration in animals showed acute inflammation with oedema and haemorrhages at the injected site.

Studies on reproductive toxicity have not been conducted.

Cu (MIBI)<sub>4</sub> BF<sub>4</sub> showed no genotoxic activity in the Ames, CHO/HPRT and sister chromatid exchange tests.

At cytotoxic concentrations, an increase in chromosome aberration was observed in the human lymphocyte assay. No genotoxic activity was observed in the in vivo mouse micronucleus test at 9 mg/kg.

Studies to assess the carcinogenic potential of the radiopharmaceutical kit have not been conducted.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Stannous chloride dihydrate  
Cysteine hydrochloride monohydrate  
Sodium Citrate  
Mannitol

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 12.

### **6.3 Shelf life**

2 years.

After reconstitution and radiolabelling, do not store above 25° C and use within 10 hours.

### **6.4 Special precautions for storage**

Do not store above 25°C. Keep the vials in the outer carton, in order to protect from light.

For storage conditions of the reconstituted after radiolabelling of the medicinal product, see section 6.3.

Storage of radiopharmaceuticals should be in accordance with national regulation on radioactive materials.

## 6.5 Nature and contents of container

15 mL multidose glass vial, type I borosilicate glass sealed with a bromobutyl rubber stopper and an aluminium caps.

Pack size: 5 vials

## 6.6 Special precautions for disposal

### General warnings

Radiopharmaceuticals should be received, used and administered only by authorised persons in designated clinical settings. Their receipt, storage, use, transfer and disposal are subject to the regulations and/or appropriate licences of the local competent authority.

Radiopharmaceuticals should be prepared in a manner which satisfies both radiation safety and pharmaceutical quality requirements. Appropriate aseptic precautions should be taken.

Contents of the vial are intended only for use in the preparation of technetium (<sup>99m</sup>Tc) sestamibi and are not to be administered directly to the patient without first undergoing the preparative procedure.

For instructions on extemporary preparation of the medicinal product before administration, see section 12.

If at any time in the preparation of this product the integrity of this vial is compromised it should not be used.

Administration procedures should be carried out in a way to minimize risk of contamination of the medicinal product and irradiation of the operators. Adequate shielding is mandatory.

The content of the kit before extemporary preparation is not radioactive. However, after sodium pertechnetate (<sup>99m</sup>Tc), is added, adequate shielding of the final preparation must be maintained.

The administration of radiopharmaceuticals creates risks for other persons from external radiation or contamination from spill of urine, vomiting or any other biological fluids. Radiation protection precautions in accordance with national regulations must therefore be taken.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## 7. MARKETING AUTHORISATION HOLDER

CIS bio international  
B.P.32  
F-91192 Gif sur -Yvette Cedex

## 8. MARKETING AUTHORISATION NUMBER

Denmark: MTn°02241  
Finland: Mtnr 25091  
Ireland: PA 0677/018/001  
Norway: MT nr 08-5992  
United Kingdom: PL 11876/0019

## 9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation:

Date of last renewal:

Denmark:	10/10/2008	12/03/2013
Finland:	09/12/2008	03/04/2013
Ireland:	12/12/2008	21/04/2013
Norway:	31/03/2009	21/04/2013
United Kingdom:	07/10/2008	24/04/2013

## 10. DATE OF REVISION OF THE TEXT

04/2024

## 11. DOSIMETRY

Technetium ( $^{99m}\text{Tc}$ ) is produced by means of a ( $^{99}\text{Mo}/^{99m}\text{Tc}$ ) generator and decays with the emission of gamma radiation with a mean energy of 140 keV and a half-life of 6.02 hours to technetium ( $^{99}\text{Tc}$ ) which, in view of its long half-life of  $2.13 \times 10^5$  years can be regarded as quasi stable.

The data listed below are from ICRP 128 and are calculated according to the following assumptions: After intravenous injection the substance is rapidly cleared from the blood and taken up predominantly in muscular tissues (including heart), liver and kidneys with a smaller amount in salivary glands and thyroid. When the substance is injected in conjunction with a stress test, there is a considerable increase of the uptake in heart and skeletal muscles, with a correspondingly lower uptake in all other organs and tissues. The substance is excreted by the liver and kidneys in the proportions 75% and 25% respectively.

Organ	Absorbed dose per unit activity administered [mGy/MBq] (resting subject)				
	Adult	15-years	10-years	5-years	1-year
Adrenals	0.0075	0.0099	0.015	0.022	0.038
Bone surfaces	0.0082	0.010	0.016	0.021	0.038
Brain	0.0052	0.0071	0.011	0.016	0.027
Breast	0.0038	0.0053	0.0071	0.011	0.020
Gall bladder wall	0.039	0.045	0.058	0.10	0.32
Gastrointestinal tract:					
Stomach wall	0.0065	0.0090	0.015	0.021	0.035
Small intestine wall	0.015	0.018	0.029	0.045	0.080
Colon wall	0.024	0.031	0.050	0.079	0.15
Upper large intestine wall	0.027	0.035	0.057	0.089	0.17
Lower large intestine wall	0.019	0.025	0.041	0.065	0.12
Heart wall	0.0063	0.0082	0.012	0.018	0.030
Kidneys	0.036	0.043	0.059	0.085	0.15
Liver	0.011	0.014	0.021	0.030	0.052
Lungs	0.0046	0.0064	0.0097	0.014	0.025
Muscles	0.0029	0.0037	0.0054	0.0076	0.014
Oesophagus	0.0041	0.0057	0.0086	0.013	0.023
Ovaries	0.0091	0.012	0.018	0.025	0.045
Pancreas	0.0077	0.010	0.016	0.024	0.039
Red marrow	0.0055	0.0071	0.011	0.030	0.044
Salivary glands	0.014	0.017	0.022	0.015	0.026
Skin	0.0031	0.0041	0.0064	0.0098	0.019
Spleen	0.0065	0.0086	0.014	0.020	0.034
Testes	0.0038	0.0050	0.0075	0.011	0.021
Thymus	0.0041	0.0057	0.0086	0.013	0.023
Thyroid	0.0053	0.0079	0.012	0.024	0.045
Urinary bladder wall	0.011	0.014	0.019	0.023	0.041
Uterus	0.0078	0.010	0.015	0.022	0.038
Remaining organs	0.0031	0.0039	0.0060	0.0088	0.016
<b>Effective dose [mSv/MBq]</b>	<b>0.0090</b>	<b>0.012</b>	<b>0.018</b>	<b>0.028</b>	<b>0.053</b>

Organ	Absorbed dose per unit activity administered [mGy/MBq] (exercise)				
	Adult	15-years	10-years	5-years	1-year
Adrenals	0.0066	0.0087	0.013	0.019	0.033
Bone surfaces	0.0078	0.0097	0.014	0.020	0.036
Brain	0.0044	0.0060	0.0093	0.014	0.023
Breast	0.0034	0.0047	0.0062	0.0097	0.018
Gall bladder wall	0.033	0.038	0.049	0.086	0.26
Gastrointestinal tract:					
Stomach wall	0.0059	0.0081	0.013	0.019	0.032
Small intestine wall	0.012	0.015	0.024	0.037	0.066
Colon wall	0.019	0.025	0.041	0.064	0.12
Upper large intestine wall	0.022	0.028	0.046	0.072	0.13
Lower large intestine wall	0.016	0.021	0.034	0.053	0.099
Heart wall	0.0072	0.0094	0.010	0.021	0.035
Kidneys	0.026	0.032	0.044	0.063	0.11
Liver	0.0092	0.012	0.018	0.025	0.044
Lungs	0.0044	0.0060	0.0087	0.013	0.023
Muscles	0.0032	0.0041	0.0060	0.0090	0.017
Oesophagus	0.0040	0.0055	0.0080	0.012	0.023
Ovaries	0.0081	0.011	0.015	0.023	0.040
Pancreas	0.0069	0.0091	0.014	0.021	0.035
Red marrow	0.0050	0.0064	0.0095	0.013	0.023
Salivary glands	0.0092	0.011	0.0015	0.0020	0.0029
Skin	0.0029	0.0037	0.0058	0.0090	0.017
Spleen	0.0058	0.0076	0.012	0.017	0.030
Testes	0.0037	0.0048	0.0071	0.011	0.020
Thymus	0.0040	0.0055	0.0080	0.012	0.023
Thyroid	0.0044	0.0064	0.0099	0.019	0.035
Urinary bladder wall	0.0098	0.013	0.017	0.021	0.038
Uterus	0.0072	0.0093	0.014	0.020	0.035
Remaining organs	0.0033	0.0043	0.0064	0.0098	0.018
<b>Effective dose [mSv/MBq]</b>	<b>0.0079</b>	<b>0.010</b>	<b>0.016</b>	<b>0.023</b>	<b>0.045</b>

The effective dose has been calculated according to a voiding frequency of 3.5 hours in adults.

#### *Cardiac imaging*

The effective dose resulting from the administration of a maximal recommended activity of 1,600 MBq of technetium ( $^{99m}\text{Tc}$ ) sestamibi for an adult weighing 70 kg is about 13.0 mSv if implementing the one-day protocol with administration of 400 MBq at rest and 1,200 MBq at exercise.

For this administered activity of 1,600 MBq the typical radiation dose to the target organ heart is 11.2 mGy and the typical radiation doses to the critical organs gall bladder, kidneys and upper large intestine are 55.2, 45.6 and 37.2 mGy, respectively.

The effective dose resulting from the administration of a maximal recommended activity of 1,200 MBq

(600 MBq at rest and 600 MBq at exercise) of technetium ( $^{99m}\text{Tc}$ ) sestamibi for a two-day protocol for an adult weighing 70 kg is about 10.1 mSv.

For this administered activity of 1,200 MBq the typical radiation dose to the target organ heart is 8.1 mGy and the typical radiation doses to the critical organs gall bladder, kidneys and upper large intestine are 43.2, 37.2 and 29.4 mGy, respectively.

#### *Scintimammography*

The effective dose resulting from the administration of a maximal recommended activity of 1,000 MBq of technetium ( $^{99m}\text{Tc}$ ) sestamibi for an adult weighing 70 kg is about 9 mSv.

For an administered activity of 1,000 MBq the typical radiation dose to the target organ breast is 3.8 mGy and the typical radiation doses to the critical organs gall bladder, kidneys and upper large intestine are 39, 36 and 27 mGy, respectively.

#### *Parathyroid imaging*

The effective dose resulting from the administration of a maximal recommended activity of 900 MBq of technetium ( $^{99m}\text{Tc}$ ) sestamibi for an adult weighing 70 kg is about 8.1 mSv.

For an administered activity of 900 MBq the typical radiation dose to the target organ thyroid is 4.8 mGy and the typical radiation doses to the critical organs gall bladder, kidneys and upper large intestine are 35.1, 32.4 and 24.3 mGy, respectively.

## **12. INSTRUCTIONS FOR PREPARATION OF RADIOPHARMACEUTICALS**

Withdrawals should be performed under aseptic conditions. The vials must not be opened before disinfecting the stopper, the solution should be withdrawn via the stopper using a single dose syringe fitted with suitable protective shielding and a disposable sterile needle or using an authorised automated application system.

If the integrity of this vial is compromised, the product should not be used.

### **Instructions for preparation of technetium ( $^{99m}\text{Tc}$ ) sestamibi**

Preparation of technetium ( $^{99m}\text{Tc}$ ) sestamibi from the kit is to be done according to the following procedure, in compliance with aseptic and radioprotection rules:

#### **Method of preparation**

##### **A. Boiling procedure**

- 1 Waterproof gloves should be worn during the preparation procedure. Remove the plastic disc from the vial and disinfect the surface of the vial closure.
- 2 Place the vial in a suitable radiation shield appropriately labelled with date, time of preparation, volume and activity.

- 3 With a sterile shielded syringe, aseptically obtain approximately 1 to 3 mL of the sterile, non-pyrogenic Sodium Pertechnetate ( $^{99m}\text{Tc}$ ) solution (200 MBq to 11.1 GBq).
- 4 Aseptically add the Sodium Pertechnetate ( $^{99m}\text{Tc}$ ) solution to the vial in the lead shield. Without withdrawing the needle, remove an equal volume of headspace to maintain atmospheric pressure within the vial.
- 5 Shake vigorously about 5 to 10 quick upside-down motions.
- 6 Remove the vial from the lead shield and place it **upright** in an appropriate boiling water bath, such that the vial is not directly in contact with the bottom of the bath and keep boiling for 10 minutes. The bath must be shielded. Timing for the 10 minutes starts as soon as the water **begins to boil** again.  
  
**Note:** The vial **must** remain upright during the boiling step. Use a water bath where the stopper will be above the level of the water.
- 7 Remove the vial from the water bath and allow to cool for 15 minutes.
- 8 Inspect visually the vial content for the absence of particulate matter and discoloration prior to administration.
- 9 Aseptically withdraw technetium ( $^{99m}\text{Tc}$ ) sestamibi using a sterile shielded syringe. Use within 10 hours of preparation.
- 10 Radiochemical purity should be checked prior to patient administration according to the radio TLC method as detailed below.

#### B. Heating block procedure

- 1 Waterproof gloves should be worn during the preparation procedure. Remove the plastic disc from the Kit vial and disinfect the surface of the vial closure.
- 2 Place the vial in a suitable radiation shield appropriately labelled with date, time of preparation, volume and activity.
- 3 With a sterile shielded syringe, aseptically obtain approximately 1 to 3 mL of the sterile, non-pyrogenic Sodium Pertechnetate ( $^{99m}\text{Tc}$ ) solution (200 MBq to 11.1 GBq).
- 4 Aseptically add the Sodium Pertechnetate ( $^{99m}\text{Tc}$ ) solution to the vial in the lead shield. Without withdrawing the needle, remove an equal volume of headspace to maintain atmospheric pressure within the vial.
- 5 Shake vigorously, about 5 to 10 quick upside-down motions.
- 6 Place the vial in the heating block previously heated to 100°C, and incubates for 15 min. The heating block should be adapted to the size of the vial in order to ensure a correct transfer of heat from the heating device to the content of the vial.
- 7 Remove the vial from the heating block and allow to cool for 15 minutes.
- 8 Inspect visually the vial content for the absence of particulate matter and discoloration prior to administration.
- 9 Aseptically withdraw technetium ( $^{99m}\text{Tc}$ ) sestamibi using a sterile shielded syringe. Use within 10 hours of preparation.

10 Radiochemical purity should be checked prior to patient administration according to the Radio TLC Method as detailed below.

### Quality Control

#### **Method**

Thin Layer Chromatography

#### **Materials**

- 1 Aluminium Oxide plate, J.T. Baker « Baker-flex » IB-FTLC , pre-cut to 2.5 cm x 7.5 cm.
- 2 Ethanol 768 g/L
- 3 Activimeter for measuring radioactivity in the 0.7 – 12 GBq range.
- 4 1 mL syringe with a 22-26 gauge needle.
- 5 Small developing tank with cover, (100 mL beaker covered with plastic film is sufficient).

#### **Procedure**

- 1 Pour enough ethanol into the developing tank (beaker) to have a depth of 3-4 mm of solvent. Cover the tank (beaker) with plastic film and allow it to equilibrate for approximately 10 minutes.
- 2 Apply 1 drop of ethanol, using a 1 mL syringe with a 22-26 gauge needle on to the Aluminium Oxide TLC plate, 1.5 cm from the bottom. Do not allow the spot to dry.
- 3 Apply 1 drop of the kit solution on top of the ethanol spot. Let the spot dry. **Do not heat.**
- 4 Develop the plate until the solvent rises to a distance of 5.0 cm from the spot.
- 5 Cut the strip 4.0 cm from the bottom, and measure the count rate of each piece in the activimeter.
- 6 Calculate the % Radiochemical purity as:

$$\% \text{ technetium } (^{99m}\text{Tc}) \text{ sestamibi} = \frac{(\text{Activity top portion})}{(\text{Total Activity})} \times 100$$

The radiochemical purity should be more than or equal to 94 %, otherwise the preparation should be discarded.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.